

Camp Creek / Walton

Aquatic plant data collected on July 27, 2000

Percent area covered with aquatic vegetation (PAC, %)	.
Percent of lake's volume filled with vegetation (PVI, %)	.
Average emergent plant biomass (kg wet wt/m ²)	3.2
Average floating-leaved plant biomass (kg wet wt/m ²)	0.7
Average submersed plant biomass (kg wet wt/m ²)	1.2
Average width of emergent and floating-leaved zone (ft.)	82.3
Average lake depth (m)	.

Frequency that plant species occur in 10 evenly spaced transects around the lake.

* Indicates non-native plant species

<u>Common Name</u>	<u>Plant Species</u>	<u>Frequency (%)</u>
black rush	<i>Juncus roemerianus</i>	80
fragrant water-lily	<i>Nymphaea odorata</i>	70
duck-potato	<i>Sagittaria lancifolia</i>	70
rush fuirena	<i>Fuirena scirpoidea</i>	60
torpedograss*	<i>Panicum repens</i>	60
sawgrass	<i>Cladium jamaicense</i>	50
slender spikerush	<i>Eleocharis baldwinii</i>	40
cone-spur bladderwort	<i>Utricularia gibba</i>	40
salt-bush	<i>Baccharis spp.</i>	30
water-pennywort	<i>Hydrocotyle umbellata</i>	30
wax myrtle	<i>Myrica cerifera</i>	30
widgeon grass	<i>Ruppia maritima</i>	30
leconte sedge	<i>Cyperus lecontei</i>	20
salt marsh fringe rush	<i>Fimbristylis castanea</i>	20
swamp rosemallow	<i>Hibiscus grandiflorus</i>	20
tapegrass	<i>Vallisneria americana</i>	20
saltmarsh false foxglove	<i>Agalinis maritima</i>	10
bacopa	<i>Bacopa monnieri</i>	10
club-rush	<i>Eleocharis cellulosa</i>	10
hatpin	<i>Eriocaulon decangulare</i>	10
large-headed rush	<i>Juncus megacephalus</i>	10
red ludwigia	<i>Ludwigia repens</i>	10
bog-moss	<i>Mayaca fluviatilis</i>	10
switchgrass	<i>Panicum virgatum</i>	10
Florida reimar grass	<i>Reimarochloa oligostachya</i>	10
seaoats	<i>Uniola paniculata</i>	10