

## Draper / Walton

Aquatic plant data collected on May 14, 2009

Percent area covered with aquatic vegetation (PAC, %)	0.0
Percent of lake's volume filled with vegetation (PVI, %)	0.0
Average emergent plant biomass (kg wet wt/m <sup>2</sup> )	1.0
Average floating-leaved plant biomass (kg wet wt/m <sup>2</sup> )	0.0
Average submersed plant biomass (kg wet wt/m <sup>2</sup> )	0.0
Average width of emergent and floating-leaved zone (ft.)	5.2
Average lake depth (m)	1.8

Frequency that plant species occur in 10 evenly spaced transects around the lake.

\* indicates non-native plant species

<u>Common Name</u>	<u>Plant Species</u>	<u>Frequency (%)</u>
black rush	<i>Juncus roemerianus</i>	90
torpedograss*	<i>Panicum repens</i>	70
seashore dropseed	<i>Sporobolus virginicus</i>	50
salt-bush	<i>Baccharis spp.</i>	30
sawgrass	<i>Cladium jamaicense</i>	20
duck-potato	<i>Sagittaria lancifolia</i>	20
smooth cord grass	<i>Spartina alterniflora</i>	20
green algae	<i>Chlorophyta</i>	10
rush fuirena	<i>Fuirena scirpoidea</i>	10
swamp rosemallow	<i>Hibiscus grandiflorus</i>	10
water-pennywort	<i>Hydrocotyle umbellata</i>	10
baby-tears	<i>Micranthemum umbrosum</i>	10
marshhay cordgrass	<i>Spartina patens</i>	10
cat-tail	<i>Typha spp.</i>	10
sea oats	<i>Uniola paniculata</i>	10